



IJCRR
Section: Healthcare
Sci. Journal
Impact Factor
4.016
ICV: 71.54

Study of Various Histopathological Patterns in Prostate Biopsy

Mital Yadav¹, Hemina Desai², Hansa Goswami³

¹3rd Year Resident, Department of Pathology, B.J. Medical College, Ahmedabad-380016; ²Associate Professor of Pathology, Department of Pathology, B.J. Medical College, Ahmedabad-380016; ³Professor and Head of Pathology, Department of Pathology, B.J. Medical College, Ahmedabad-380016.

ABSTRACT

Introduction: Diseases of prostate gland are important source of morbidity and mortality in male patients. The spectrum of diseases consists of inflammatory conditions, nodular hyperplasia, malignancy etc. The risk of diseases increases with age. Prostate biopsy is essentially a test which detects cancer and other benign conditions of the prostate in patients who have specific indications for it. Other modalities of diagnosis also have been developed, comprising of radiological modalities and prostatic biomarkers. However, biopsy with histopathological examination is the gold standard for the final diagnosis.

Aims and Objective: To study various histopathological patterns in Transurethral resection of prostate (TURP) and needle biopsy of prostate, including benign prostatic hyperplasia (BPH), prostatitis (identifying type and grade of inflammation), incidental detection of carcinoma prostate and confirmation of diagnosis of clinically suspected carcinoma with application of modified Gleason system for grading.

Materials and Methods: Total 100 cases of prostatic lesions including transurethral resection of prostate (TURP) and needle biopsy were evaluated in this study. The received specimens were fixed in 10% neutral buffered formalin and routine paraffin processing followed by hematoxyline and eosin staining was done. All the specimens were analysed with following parameters: type of specimen, age of patient, histopathological pattern and final diagnosis.

Results: In total 100 cases of present study, the most common age group presenting prostatic lesions was 61-70 years. The youngest patient presenting with prostatic lesion was 45 years age, and oldest was of 85 years of age. In this, various histopathological patterns observed were- Benign prostatic hyperplasia (BPH) in 45%, benign prostatic hyperplasia (BPH) with co-existing chronic prostatitis in 10%, granulomatous prostatitis 1% (single case), acute prostatitis 3% and acute on chronic prostatitis 6%. Less frequent finding were benign prostatic hyperplasia (BPH) with basal cell hyperplasia 1%, benign prostatic hyperplasia (BPH) with squamous metaplasia 2%, benign prostatic hyperplasia (BPH) with clear cell changes 3% and some of biopsy show only stromal tissue with hyperplasia in 4%. We reported 7 cases of adenocarcinoma prostate with application of modified Gleason grading system.

Conclusion: Benign lesions are more common than malignant lesions. Among the histological patterns of prostatic lesions, benign prostatic hyperplasia (BPH) is predominant type. It is necessary to study all prostate biopsy [Transurethral resection of prostate (TURP) and needle] in order to identify premalignant lesions, proliferative activity, and grade of inflammation. Histopathological diagnosis and grading plays a definitive role in the management of prostatic cancer.

Key Words: Transurethral resection of prostate (TURP) and Needle biopsy, Benign prostatic hyperplasia (BPH), Prostate Carcinoma

INTRODUCTION

The prostate gland is the pear shaped glandular organ that weighs up to 20 gms¹. Prostate is fibromusculoglandular organ encircling the neck of the urinary bladder. The prostate is an exocrine gland and forms a significant component of seminal fluid. Histologically it consists of glands lined by basal cuboidal cells and inner secretory columnar cells (double

layered)¹. Most patients of benign lesions present with complaints related to micturition and incontinence. Of the diseases which affects the prostate the most frequently encountered in clinical practice are benign prostatic hyperplasia (BPH), prostatic cancer and prostatitis².

Benign prostatic hyperplasia is an extremely common condition in men over the age of 50 years and shows remarkable

Corresponding Author:

Dr. Mital Yadav, 3rd Year Resident, Department of Pathology, B.J. Medical College, Ahmedabad-380016,
Mobile: 9924029075; E-mail: mitalyadav1991@gmail.com

ISSN: 2231-2196 (Print)

ISSN: 0975-5241 (Online)

DOI: 10.7324/IJCRR.2017.9219

Received: 28.09.2017

Revised: 12.10.2017

Accepted: 27.10.2017

racial and geographical variations in incidence and mortality³. The clinical incidence of this disease is only 8% during the 4th decade, but it reaches 50% in the 5th decade and 75% in the 8th decade¹.

Benign prostatic hyperplasia (BPH) is not premalignant lesion for the prostatic cancer but it may be related to prostate cancer arising in transition zone (Difenbach et al., 2002)⁴. Non-specific granulomatous prostatitis is noticed occasionally in prostate specimen. It was first described by Tanner and Mc Donald in 1943, who reported an incidence of 3.3% of granulomatous prostatitis in inflammatory lesions. Recently premalignant lesions have become defined, largely as a result of advances in technology.

Carcinoma of prostate is most common internal malignancy among men in United State and is responsible for 10% of cancer deaths⁵. In India, prostate cancers constitute about 5% of all cancers in male^{5,6}. Screening of prostatic lesions constitute prostate specific antigen, digital rectal examination, and transrectal ultrasound, but biopsy remains the gold standard diagnostic tool for final diagnosis. Prior to the PSA era, up to 27% of prostate cancers were detected incidentally at the time of transurethral resection of prostate (TURP)⁷. The modified Gleason's system appears better to predict progression –free survival after radical prostatectomy than the original Gleason's system did.

MATERIAL AND METHODS

The study was conducted in the Department of Pathology, B.J Medical College, civil hospital – a tertiary level hospital of Ahmedabad. The study period was from April 2017 to June 2017 and prostatic specimen data were collected retrospectively from archives of histopathology register. From April 2017 to June 2017 a total of 100 transurethral resection of prostate (TURP) and needle biopsy were evaluated. The received specimens were fixed in 10% neutral buffered formalin and routine paraffin processing followed by hematoxyline and eosin staining was done. All the specimens were analysed with following parameters: type of specimen, age of patient, histopathological pattern and final diagnosis. The slides were examined under light microscope, and Various histopathological patterns were studied in all cases and were classified with reference to age. Following histopathologic assessment, the tumors were classified according to WHO recommendation, and histologic grading was done using modified Gleason's system.

RESULT

In this study a total of 100 prostate specimens were studied which were received between April 2017 to June 2017

in the histopathology section of department of Pathology in B.J. Medical College, Civil hospital, Ahmedabad. Of the 100 specimens 47(47%) were transurethral resection of prostate (TURP) specimens and 53(53%) needle biopsy specimens.

All prostatic specimens were broadly classified into benign 93(93%) and malignant 7(7%) [figure 1]. Maximum cases of benign prostatic hyperplasia (BPH) (38) were seen in the 61-70 years age group [Table 1 and figure 2]. A glandulostromal pattern of hyperplasia (72%) was the most frequent histological pattern which occurred. Stromal pattern of hyperplasia (4%) was less common. In this 45(45%) were benign prostatic hyperplasia (BPH) only [Figure 3]. Benign prostatic hyperplasia (BPH) with co-existing chronic prostatitis 10(10%) [Figure 4], acute on chronic prostatitis 6(6%), acute prostatitis 3(3%) [Figure 5] and granulomatous prostatitis 1(1%) [Table 2]. In this study we reported one (1%) case of granulomatous prostatitis which showed granulomas with giant cells; however, Ziehl Nelsen staining was negative for AFB, and was labeled as non specific.

Less frequent finding was benign prostatic hyperplasia (BPH) with clear cell cribriform hyperplasia 3(3%) [Figure 6], benign prostatic hyperplasia (BPH) with basal cell hyperplasia 1(1%) [figure 7], benign prostatic hyperplasia (BPH) with squamous metaplasia 2(2%) [Figure 8].

We reported 7 cases of adenocarcinoma prostate with modified Gleason score. Histopathology of biopsy received showed single, separate, much more variable glands, closely packed but irregularly separated, ragged and poorly defined edge (Gleason's grade 3) [Figure 9]. Second most predominating pattern is fused glandular, cribriform pattern and hypernephroid (Gleason's grade 4) [Figure 10]. The most common predominant grades observed in this study were grade 3 and grade 4. The most common score obtained was 6 in 4 cases out of the total 7 adenocarcinoma cases.

DISCUSSION

A retrospective study of 100 cases of prostate biopsy were done, result were noted and comparison with other studies were undertaken as follows:

Age

Lesions of prostate are extremely common over the age of 50 years. The clinical incidence of this disease is only 8% during the 4th decade, but it reaches 50% in the 5th decade and 75% in the 8th decade¹. Benign prostatic hyperplasia (BPH) and carcinoma of the prostate are increasingly frequent with advancing age⁸.

In present study, age group affected with prostatic pathology was 45-85 years. Maximum cases of benign prostatic

hyperplasia (BPH) (38) were seen in the 61-70 years age group similar to Matapurkar et al⁹. Kim KB et al. had more cases(44) in 71-80 age group¹⁰. Malignant lesions encountered predominantly in age group 61-70 years that are similar to Sharma et al¹¹.

Benign and malignant

Benign prostatic hyperplasia (BPH) and adenocarcinoma are the two most common conditions affecting prostate gland.

In present study malignant cases were 7 (7%), and benign cases were 93 (93%) with ratio 1:14 approximately. In present study, the predominant lesion was benign prostatic hyperplasia (BPH) (45%) similar to Neha Angurana's¹² study (50.5%), followed by benign prostatic hyperplasia (BPH) with prostatitis (21%) and prostate cancer (7%), basal cell hyperplasia and cribriform clear cell hyperplasia was observed.

Microscopy and Histopathological patterns

Benign prostatic hyperplasia: There is hyperplasia of glandular and stromal tissue with papillary buds, in foldings and cysts. In glandulostromal pattern-Proliferating glands and fibromuscular stroma were assessed to be roughly in equal proportion. While in Stromal pattern- The sections showed more stromal elements than glands or were made up almost entirely of stromal elements.

In present study, glandulostromal pattern of hyperplasia(72%)was the most frequent histological pattern which occurred. Stromal pattern of hyperplasia (4%)was less common. Our study was in agreement with study done by Zeenath Begum, Abdul Hakeem Attar, Mandakini B. Tengli, Mohammed Mateen Ahmed¹³ (8%) in respect that glandulostromal hyperplasia was most commonly noted.

Prostatitis: It shows inflammatory change with in prostate glands. Acute prostatitis:where there are neutrophils within acini and throughout stroma, intraductal desquamated cellular debris, stromal edema and hyperaemia. Acute prostatitis is rarely seen in surgical specimens. In present study acute prostatitis(3%)was observed. Chronic prostatitis:Chronic prostatitis is more common. Chronic inflammation typically involves the prostate in periglandular distribution and contains an admixture of plasma cells,lymphocytes and histiocytes. In present study Benign prostatic hyperplasia (BPH) with co-existing chronic prostatitis in 10(10%) cases was observed. Chronic prostatitis is most commonly observed in benign prostatic hyperplasia (BPH)which was also observed by Dr. Ashish Joshee, Dr. Kaushal C.L. Sharma (14) in their study¹⁴. Malakoplakia:Usually associated with bladder disease, may clinically mimic cancer, resulting in prostatic induration and a hypoechoic lesion seen on transrectal ultrasound. On microscopy it show Marked histiocytic infiltrate containing Michaelis-Gutmann bodies (calcospherites) that

may resemble signet ring cells at frozen section¹⁵.

Granulomatous prostatitis: Most commonly diagnosed granulomatous process within the prostate is non specific. It is thought to represent an initially immune-mediated process accompanied by a reaction to the prostatic secretions released from obstructed ducts on microscopic examination-large nodular aggregate of histiocytes, epithelioid cells, lymphocytes, and plasma cells. Characteristically, granuloma formation are seen¹⁶.

In present study granulomatous prostatitis(1%) was observed. The incidence of non specific granulomatous prostatitis was 0.5% observed in Stillwell TJ, Engen DE, Farrow GM¹⁷ in their study.

Types of epithelial hyperplasia

Various hyperplastic lesions were observed like papillary, basal, cribriform. basal cell hyperplasia is usually seen in the transitional zone, but it may also occur in the peripheral portion of the gland. Microscopically it appears as small, solid nests of benign-appearing epithelial cells with a somewhat clear cytoplasm¹. Cribriform hyperplasia is an uncommon and controversial form of hyperplasia¹. In clear cell cribriform hyperplasia, cytoplasm of hyperplastic glandular cells often a clear¹.

In present study benign prostatic hyperplasia (BPH)with basal cell hyperplasia 1(1%) and clear cell cribriform hyperplasia 3 (3%)was observed.

Types of metaplasia

Squamous metaplasia can be seen at the periphery of infarcts, after TUR, as a result of hormonal manipulation, or sometimes with no obvious predisposing cause. In present study benign prostatic hyperplasia (BPH) with squamous metaplasia in 2(2%) cases was observed.

Carcinoma Prostate

Carcinoma of prostate is the second most common malignancy in men, second only to lung cancers. Hormonal factors play important role in the development of prostatic carcinoma.5-10% of prostatic carcinoma have a genetic link. A higher incidence has been seen in males with a first degree relative being affected¹.

Features favoring prostatic carcinoma

Architectural: Infiltrative patterns, small glands, crowded glands.

Nuclear: Prominent nucleoli, enlargement, hyperchromasia, mitotic figures, apoptotic bodies.

Cytoplasmic: Amphophilic cytoplasm, sharp luminal border¹⁸.

In present study carcinoma prostate was identified incidentally in 7 (7%) i.e., 7 of 100 cases, close to previous study by Otto B et al.¹⁹ where only 1.4% patients were found to have prostate cancer on pathology in prostate biopsy samples. Adenocarcinoma was the only histological type of cancer found in this study. Adenocarcinomas were graded according to Gleason's system by taking into account morphological appearance of glandular cells and the glandular pattern. The most common predominant grades observed in this study were grade 3 and grade 4. The most common score obtained was 6 in 4 cases out of 7 adenocarcinoma cases. Vollmer²⁰ had score 6 most common whereas Brawn et al.²¹ had scores 6 and 7 in his study.

CONCLUSION


We conclude that prostatic lesions are common in age group of 61-70 years. Benign conditions are more common than malignant conditions. Among the histological types of prostatic lesions, benign prostatic hyperplasia (BPH) is predominant type, followed by benign prostatic hyperplasia (BPH) with prostatitis. It is necessary to reassess periodically all prostate biopsies carefully in order to identify premalignant lesions, proliferative activity, grade of inflammation. Efforts should be made to apply modified Gleason's system in case of adenocarcinoma of prostate to improve management.

REFERENCES




1. Rosai and Ackerman's surgical pathology tenth edition.
2. Cotran RS, Kumar V, Robbins SI. Prostate. In: Cotran RS, Kumar V, Robbins SI (eds): Robbins Pathologic Basis Of Disease, 6th ed., Philadelphia: Saunders Co., pp. 1025-1034, 1994.
3. Walsh PC. Benign prostatic hyperplasia. In: Walsh PC, Gittes RF, Perlmutter AD, Stamey TA (eds): Campbell's Urology 5th ed., vol.2, Philadelphia: W.B. Saunders Co., pp. 1248-1265, 1986.
4. Difenbach MA, Dorsey J, Uzzo RG, Hanks GE, Greenberg RE and Horwitz E *et al.*, (2002). Decision making strategies for patients with localized prostate cancer, Seminars in Urologic Oncology 20:52-62.
5. Egevad L. Consolidated Report of Population Based Cancer Registries 2001-2004: Incidence and Distribution of Cancer. Bangalore (IND): Coordinating Unit, National Cancer Registry Programme, Indian Council of Medical Research; 2006.
6. Garg M, Kaur G, Malhotra V, Garg R. Histopathological spectrum of 364 prostatic specimens including immunohistochemistry with special reference to grey zone lesions. Prostate Int 2013; vol 1 (4): 146-151.
7. B. Tombal, L. de Visccher, J. P. Cosyn *et al.*, Assessing the risk of unsuspected prostate cancer in patient with Benign Prostatic hypertrophy: a 13 years retrospective study of incidence and natural history of T1a-T1b Prostate cancers, "BJU International, Vol. 84, no. p. 1015-1020, 1999.
8. A Josephine -2014, clinicopathological study of prostate biopsy, © 2014 Journal of Clinical and Diagnostic Research.
9. Matapurkar BG, Taneja OP. Incidence of carcinoma prostate. Ind J of Cancer 1969 Sept 172-182.
10. Kim KB, Kim KS. A Histopathological observation on 48 cases of benign prostatic hypertrophy. Korean J Urol 1982; 23(8):30.
11. Sharma GC, Mathur SC, Sharma ML. Occult carcinoma in benign hypertrophy of prostate (Clinicopathological study of 100 cases). Ind J Surg 1972 April: 152-155.
12. Neha angurana, pattern of prostate diseases- a histopathological study in Jammu, international journal of basic and applied medical sciences issn: 2277-2103 (online)
13. Zeenath Begum, Abdul Hakeem Attar, Mandakini B. Tengli, Mohammed Mateen Ahmed, Study of Various Histopathological Patterns in Turp Specimens and Incidental Detection of Carcinoma Prostate, Indian Journal of Pathology and Oncology, October – December 2015;2(4):303-308.
14. Dr. Ashish Joshee, Dr. Kaushal C.L. Sharma; The histomorphological study of prostate lesions, IOSR Journal of Dental and Medical Sciences (IOSR-JDMS) e-ISSN:
15. Wagner D, Joseph J, Huang J, Xu H. Malakoplakia of prostate on needle core biopsy, a case report and review of the literature.
16. Oppenheimer JR, Kahane, Epstein JI. Granulomatous prostatitis on needle biopsy, Arch pathol Lab Med 1997.
17. Stillwell TJ, Engen DE, Farrow GM. The clinical spectrum of granulomatous prostatitis: A report of 200 cases. J Urol. 1987; 138:320-323.
18. Epstein, George J, Netto. Biopsy interpretation of the prostate, 4th edition.
19. Otto B *et al.* incidental prostate cancer in Transurethral Resection of the Prostate Specimen in moderate era. Advances in Urology 2014; 4:1-4.
20. Vollmer RT. Prostatic cancer and chip specimen: complete versus partial sampling. Hum Pathol 1986; 17:285-290.
18. Otto B *et al.* incidental prostate cancer in Transurethral Resection of the Prostate Specimen in moderate era. Advances in Urology 2014; 4:1-4.
21. Brawn PN, Ayala AG, vonEschenbach AC, Hussey DH, Johnson DE. Histologic grading study of prostate adenocarcinoma: The development of a new system and comparison with other methods- A preliminary study. Cancer 1982; 49:525-532.

Table 1: Age-wise distribution of cases:

Age (yr)	Benign,n(%)	Malignant, n(%)
41-50	6	0
51-60	24	1
61-70	38	4
71-80	23	2
81-90	2	0
Total	93	7

**Table 2: Distribution of Histopathological Patterns pattern:**

Histopathological patterns	No. of cases	Percentage
Benign prostatic hyperplasia (BPH)alone	45	45%
Benign prostatic hyperplasia (BPH)with chronic prostatitis	10	10%
Benign prostatic hyperplasia (BPH)with acute on chronic prostatitis	6	6%
Granulomatous prostatitis	1	1%
Benign prostatic hyperplasia (BPH)with acute prostatitis	3	3%
Stromal hyperpasia only	4	4%
Benign prostatic hyperplasia (BPH)with Clear cell hyperplasia	3	3%
Benign prostatic hyperplasia (BPH)with basal cell hyperplasia	1	1%
Benign prostatic hyperplasia (BPH)with squamous metaplasia	2	2%
Benign prostatic tissue	18	18%
Adenocarcinoma prostate	7	7%
Total	100	100%

Figure 2: Age-wise distribution of cases.**Figure 3: Benign prostatic hyperplasia (HandE stain) (10X Magnification).****Figure 4: Chronic prostatitis (HandE stain) (20X Magnification).****Figure 1: Distribution of benign and malignant lesion.**




Figure 5: Acute prostatitis (HandE stain) hyperplasia (HandE stain) (20X Magnification).




Figure 8: Squamous metaplasia (HandE stain) (20X Magnification).




Figure 6: Clear cell cribriform hyperplasia (HandE stain) (20X Magnification).




Figure 9: Adenocarcinoma of prostate Gleason's grade (3+3). (HandE stain) (10X Magnification).




Figure 7: Basal cell hyperplasia (HandE stain) (20X Magnification).




Figure 10: Adenocarcinoma of prostate Gleason's grade (4+4). (HandE stain) (10X Magnification).